Description of Taboos Behavior Practice among Pregnant Women in West Java INDONESIA

Mira Trisyani Faculty of Nursing Universitas Padjadjaran

Outline of Presentation

- Background
- ♦ Literature Review
- Aim & Objectives
- Methodology and Method
- Data Collection
- Data Analysis
- Ethics
- Results

Purpose of Study

♦ To investigate practically of beliefs about food taboos, activity taboos and attended of special activity during pregnancy.

Background

Why recognition of taboos behavior?

- In Indonesia as generally pregnancy is considered normal phenomena
- The experience of pregnancy encompasses physiological, spiritual, and socio-cultural dimensions
- Food choices during pregnancy and lactation are influenced by a multifactor ethnics



Indonesia women assumed that pregnant women should have attention in healthy self-care and babies by: (1) concern and believe about taboos food; (2) concern with taboos activity; and (3) attended of special activities.

Background

- ▲ Infant Mortality Rate (IMR) in Indonesia was last reported at 27.20 per 1,000 live births according to World Bank report release in 2011, that was decreased since year between 1998 and 2002 (35 per 1,000 live births).
- ♦ Findings prevalence of chronic energy deficiency among pregnant women in west Java
- Differences of maternal health care practice among pregnant women

Literature Review

- Belief and culture on pregnancy:
 - Maternal Health Care
 - Focus on cultural awareness influences on nutrition

Literature Review

- ♦ Culture is a set of learned customs, beliefs, values, ways of communication, actions, and institutions shared by racial, ethnic, religious or social group that shape their manner of living.
- Since of fetus development and childbirth process is a natural phenomenon in continuity of human life, but the various groups of public with its culture in the whole world have multifarious perception, interpretation and behavior response in implications to health.

Literature Review

- ▶ The healthy babies related with health behaviors of pregnant woman include birth weight are adequate prenatal care, smoking, moderate to heavy alcohol use, and poor nutrition of the mother (Tiedje, Kingry & Stommel, 1992).
- ▶ The traditional background sometimes influences of the woman behavior during pregnancy, many beliefs and practices relating to pregnancy and childbirth among women of non-Western backgrounds have received much attention in the past few decades. (Liamputtong Rice, 2000).

Aim

• To investigate practically of beliefs about food taboos, activity taboos and attended of special activity during pregnancy.

Objectives

- 1. Identify among pregnant women of perspective in taboos food and taboos activity during pregnancy
- 2. Identify among pregnant women of the important things of attended in special activity during pregnancy

Contribution

♦ This study will contribute to knowledge about nursing care practice in Indonesia and in particular issues encountered by Indonesian health professionals in relation to maternal health care

Methodology

- ♦ The research was used descriptive quantitative design to identify the pregnant women activity especially about taboos behavior during her pregnancy in several cities in West Java.
- ◆ The instruments divide into 2 parts are: part I Demographic Data consisted of 7 items constructed by researcher; and part II Taboos Activity Questionnaires consist of 17 items, there were an instrument utilized for this study.

Research Setting and Participant

- ♦ The setting for this study was carried out at four public hospitals and one private obstetric practice of midwives from 30 December 2008 to 30 January 2009.
- ▲ A total of 428 packages of questionnaires was distribution to pregnant women but 300 pregnant women were completed the questionnaires and returned into the researcher.

Research Setting and Participant

- ♦ The criteria for selection of participants were:
 - (1) all participants were able to read, write, and understand Indonesia language.
 - (2) women who cultural background is Sundanese;
 - (3) participants were eighteen to forty-five years of age;
 - (4) participants who had pregnancy in first, second and third trimester without undue complication or any high risk;
 - (5) participants included primigravida (first birth) and multigravida (second or subsequent birth);
 - (6) All pregnant women who met the inclusion criteria and consented to participate after reading or being read the informed consent were enrolled.

Data Collection

- Quantitative data collection methods will be used for this study
- A cover letter form given to the participant explained about the purpose of the study and instructions how to complete the questionnaires.
- The form asked for decision to participant if they would to participant in this research or not.
- ▶ The questionnaires were given to each participant with the instructions of how to complete it.

Data Analysis

- Descriptive Statistics
 - Descriptive statistics will include frequency distributions, percentages

Ethics

- ♦ Ethics approval from Universitas Padjadjaran Human Research Ethics Committee has been granted
- Ethic approval from the hospital concerned has been approved
- ♦ The study will follow to procedures such as distribution of study information sheets, informed and signed consent and confidentiality of participants

Results: Demographic data

- ♦ All the participants almost a pregnant women, the ranged of age 20 to 29 years old (56%).
- Forty-six percent (46%) were pregnant women with gestational age in trimester III.
- The data showed (47%) were in multigravida (second or subsequent birth), and eighty-one percent (81%) of participants are low-educated.
- ▲ Also (74%) almost of respondent did not work. Forty-seven percent (47%) of pregnant women reported that their monthly incomes about 100 USD or less.

Results: Food and Activities Taboos

The highest mean for question "avoid to seeing a moon eclipse" (3.38± .38), it is describes that some pregnant women forbidden to see moon eclipse (83.7%). Furthermore the lowest score is the question by "avoid to eat some seafood" (2.63±.79). Other responses from participant as food taboos was "restriction to eat some kind of fruits and vegetable" (2.29±.04), although most participant answered in "rarely" (37.7%) nevertheless that was presumption needed correction.

Results

♦ The findings showed that 83.7% (n=251) of pregnant women forbid to see moon eclipse. Some pregnant women of 69.3% (n=208) keep away from unpleasant behavior during pregnancy. 37% of participants assume certain activity as a food taboos. Those are "restriction to eat some kind of fruits and vegetable like pineapple, avocado, pomegranate, guava, orange squash, durian, jack fruit, papayas, sugar cane, and eggplant".

Conclussion

♦ The traditional background sometimes influences woman behavior during pregnancy. The findings were supported by study at four regencies of West Java, Indonesia which have strongly influenced by taboos behavior. Additional studies of practically maternal health care activity among pregnant women are needed. Regarding taboos behavior, nurse should offer health education among pregnant women on proper practice maternal health care activities.

Thank you for attention and please coming to Indonesia



Ahn M. O. (2008). Pregnancy and Culture Norm: Korean Experience Presentation Detail, from http://mwia.regional.org.au/papers/session/o/speakers_oral.htm

- [2] Backstrand, J. R., Goodman A. H., Allen, L. H., Pelto, & G. H. (2004). Pulque intake during pregnancy and lactation in rural Mexico: alcohol and child growth from 1 to 57 months. Eur J. Clin. Nutr, 58(1), 626-1634
- [3] Betz, H., G. (1992). "Postmodernism and the New Middle Class" in Theory, Culture & Society Journal, Vol. 9, No. 2, 93-114
- [4] Bernhardt, J. H. (1990). Potential workplace hazards to reproductive health. Journal of Obstetric, Gynecologic, and Neonatal Nursing, 19(1),53-62
- [5] Bustan, M. N., and Coker, A. L. (1994). Maternal attitude toward pregnancy and the risk of neonatal death, *American Journal of Public Health*, 84(3), 411-414
- [6] Burk, Martha E., Wieser, Peggy C, & Keegan, Lynn. (1995). *Cultural beliefs and health behaviors of pregnant Mexican-American women:* Implications for primary care. Aspen Publishers, Inc.: Advance the Nursing Science.
- [7] Bobak, I. M., Lowdermilk, D. L., & Jensen, M. D.(1995). Maternity Nursing. Saint Louis: C.V. Mosby Company
- [8] Bulletin of World Health Organization. (2007). Programmes and Projects. 85. October.
- [9] Cao, A. Rosatelli, M.C., Monni G., & Galanello, R. (2002). Screening for thalassemia:a model of success. *Obstet Gynecol Clin North Am* 29(2), 305–328
- [10] Chapman, J. S. (1975). The Relation Between Auditory Stimulation of Short Gestation Infants and Their Gross Motor Limb Activity. Doctoral Dissertation, New York University

- [11] Chang, S. B., Park, Y. J., Choi, Y. S., & Chung, C. W. (1996). Factors of the "Taegyo" of Korean Pregnant Women: Self Care of Pregnant Women Based on Oriental Folk Behavior. J. Korean Acad. Nurs, 26(2), 345-358.
- [12] Chamberlain, D. B. (1998). Prenatal Receptivity and Intelligence. *J. prenatal and perinatal Psychology and Health*, from http://www.smartstartnc.org/conference/Handouts08/253b.doc
- [13] DeDee, L. S., PhD. (1998). Relationships among diet, nutrition beliefs, Maternal attachment and adjustment to pregnancy in the second trimester. Doctoral Dissertation, Winconsin University.
- [14] Director of West Java Regional Tourism Office. (1995). *The link of emerald of Indonesia West Java*. Official Publication West Java Government Tourist Office Indonesia.
- [15] Department of Health Republic Indonesia. (2007). Health Profile 2005. Jakarta: Department of Health.
- [16] Delaney, C. (1991). The Seed and the Soil: Gender and Cosmology in a Turkish Village Society. Berkeley: University of California Press.
- [17] Ebomoyi, E. (1988). Nutritional beliefs among rural Nigerian mothers. Ecol. Food Nutr. 22:43-52
- [18] Fishbein, E. G., et al. (1990). How safe is exercise during pregnancy? Journal of Obstetric, Gynecology, and Neonatal Nursing, 19, 45.
- [19] Frankel, B. (1977). Childbirth in the ghetto. R&E Research, San Francisco, Calif.
- [20] Harrison, G. G., Zaghloul S. S., Galal O. M., & Gabr, A. (1993). Breastfeeding and weaning in poor urban neighborhood in Cairo, Egypt: maternal beliefs and perceptions. *Soc Sci Med. 36*:1063-1069

- [21] Hatch, M. C., et al. (1993). Maternal exercise during pregnancy, physical fitness, and fetal growth. *American Journal of Epidemiology*, 137, 1105
- [22] Helen, H., Gifft, Marjorie, B., Washbon, Gail, G., & Harrison. (1972). Nutrition Behavior and Change. New Jersey: Prentice-hall, inc., Englewood cliffs.
- [23] Herman, B., M. D., & Perry, S. K., Ph.D. (1996). The *Twelve-Month Pregnancy. Update edition*. Los Angeles: Lowell House.
- [24] Howard, J., & Berbigilia, V. (1997). "Caring for childbearing Korean women." *Journal of Obstetrics, Gynecological, and Neonatal Nursing*, 26 (6), 665-671.
- [25] Indonesia let's go archipelago. West Java, from Http://www.Indonesia-tourism.
- [26] Julie, M. (2005). Trancultural index.
- [27] Kaplan, F. (1998). Tay–Sachs disease carrier screening: a model for prevention of genetic disease. *Genet Test 2*(4):271–292

- [28] Kent, H., Johnson, K., Curtis M., Richardson, H. J., & Atrash, H. (2006). Proceedings of the Preconception Health and Health Care Clinical, Public Health, and Consumer Workgroup Meetings Hosted by the Centers for Disease Control and Prevention, National Center on Birth Defects and Developmental Disabilities. Atlanta, Georgia.
- [29] Kim, H. J. (2008). Women between Cultures: Experiences in Pregnancy and Postpartum Care. Fieldworking online A community for Researcher and Writers.
- [30] Korenbrot, C.C., Steinberg, A., Bender, C., & Newberry, S. (2002) Preconception care: a systematic review. Matern Child Health J, 6(2),75-88
- [31] Kogan, M. D., Alexander, G. R., Kotelchuck, M., Nagey, D.A., & Jack, B.W.(1994). Comparing mother's report on the content of prenatal care received with recommended national guidelines for care. *Public Health Reports*. 109,637-646.
- [32] Kozier, B., Erb, G., Blais, K., & Wilkinson, J. M. (1995). Fundamentals of Nursing: Concepts, Process, and Practice/Clinical Companion. (5th ed.) Addison-Wesley.
- [33] Kenneth, H.H., & Wong, E.Y.A. (2007). Early Conceptus Growth and Immunobiologic Adaptation of Pregnancy. Handbook of Clinical Obstetrics: Fetus & Mother 3th edition. Blackwell Publishing Ltd.
- [34] Liamputtong, P. (2005) Pregnancy, childbirth and traditional beliefs and practices in Northern Thailand. In H. Selin (ed.), Childbirth across cultures: Ideas of pregnancy, childbirth and the postpartum period in many of the world's cultures. Springer: Dordrecht, The Netherlands.
- [35] Laffrey, S. C. (1986). Development of Health Conception Scale. Research in Nursing and Health, 9:107-113

- [36] Laderman, C. (1987). Wives and Midwives: Childbirth and Nutrition in Rural Malaysia. Berkeley: University of California Press.
- [37] Lee, K. H., Lee, Y. S., & Hee, K. J. (2004). Women's Health Nursing. Korea: Hyonmunsa.
- [38] Lorio, C. K. D. (2005). Measurement in Health Behavior: Methods for Reasearch and Education. San Fransisco: Jossey Bass, 60.
- [39] Lowdermilk, D. Perry, S., & Bobak, I. (1999). *Maternity Nursing (5tf ed.)*. Mosby: St. Louise.
- [40] Lipetz, M., & Newcomb, J. (1986). Concept Paper: Nursing Practice in Women's Health. Nursing Research, 35(2), 143

- Mardiati, N., Purnawan, J., Prastuti, C., & Purwa, K. S. (2004). Economic Analysis of the Village Midwives (BDD) Program. A Case Study on Sustainability of the Midwives in 3 districts in Java. Center for Health Research, University of Indonesia.
- [42] Mennella, J.A, Turnbull, B., Zielgler P. J, & Martinez, H. (2005). Infant feeding practices and early flavor experiences in Mexican infants: an intra-cultural study. *J. Am Diet Assoc.*, 105, 908-915
- [43] Meutia, F. S. (1998). *Kehamilan, Kelahiran, Perawatan Ibu dan Bayi dalam Konteks Budaya*. Jakarta: Universitas Indonesia
- [44] Montagu, A. (1978). Touching: The Human Significance of the Skin. New York: Harper & Row.
- [45] Muchowski, K., & Paladine, H. (2004). An ounce of prevention: the evidence supporting periconception health care. *J. Fam Pract* 53(2), 126–133
- [46] Newman, L. F. (1995). Women's Medicine: a Cross-Cultural Study of Ingenous Fertility Regulation. New Brunswick: Rutgers University Press (with helped James M. Nyce)
- [47] New nutrition map of Indonesia highlights widespread infant mortality, underweight children. Folklore, Food, Female reproductive cycle. *Ecol Food Nutr.* 7:41-49. from Http://www.Relief web.net Snow LF, Johnson SM. (1978).